



**Pawnee Montane Skipper**  
(*Hesperia leonardus montana*)



Photo Source: U.S. Department of Agriculture



03-0402-11



**Pawnee Montane Skipper**  
(*Hesperia leonardus montana*)



Photo Source: U.S. Department of Agriculture



03-0402-11



**Pawnee Montane Skipper**  
(*Hesperia leonardus montana*)



Photo Source: U.S. Department of Agriculture



03-0402-11



**Pawnee Montane Skipper**  
(*Hesperia leonardus montana*)



Photo Source: U.S. Department of Agriculture



03-0402-11

**Pawnee Montane Skipper  
(Hesperia leonardus montana)**

Federal status: Threatened  
Colorado status: Not listed

The upper side of this butterfly's wing is brownish red with distinct yellowish spots near the outer margins. The underside is brown with cream spots. The wing-span is from 1.5 to 2 inches.

Habitat: This species is found in dry, open, Ponderosa pine woodlands at elevations from 6,000 to 7,500 feet. The slopes are moderately steep with soils derived from Pikes Peak granite. The understory is limited in the pine woodlands. Blue Grama grass is the larval food plant, and the prairie gayfeather is the primary nectar plant. The species is found in Douglas, Jefferson, Park and Teller counties. The total known habitat within the range is estimated to be 37.9 square miles.



**Pawnee Montane Skipper  
(Hesperia leonardus montana)**

Federal status: Threatened  
Colorado status: Not listed

The upper side of this butterfly's wing is brownish red with distinct yellowish spots near the outer margins. The underside is brown with cream spots. The wing-span is from 1.5 to 2 inches.

Habitat: This species is found in dry, open, Ponderosa pine woodlands at elevations from 6,000 to 7,500 feet. The slopes are moderately steep with soils derived from Pikes Peak granite. The understory is limited in the pine woodlands. Blue Grama grass is the larval food plant, and the prairie gayfeather is the primary nectar plant. The species is found in Douglas, Jefferson, Park and Teller counties. The total known habitat within the range is estimated to be 37.9 square miles.



**Pawnee Montane Skipper  
(Hesperia leonardus montana)**

Federal status: Threatened  
Colorado status: Not listed

The upper side of this butterfly's wing is brownish red with distinct yellowish spots near the outer margins. The underside is brown with cream spots. The wing-span is from 1.5 to 2 inches.

Habitat: This species is found in dry, open, Ponderosa pine woodlands at elevations from 6,000 to 7,500 feet. The slopes are moderately steep with soils derived from Pikes Peak granite. The understory is limited in the pine woodlands. Blue Grama grass is the larval food plant, and the prairie gayfeather is the primary nectar plant. The species is found in Douglas, Jefferson, Park and Teller counties. The total known habitat within the range is estimated to be 37.9 square miles.



**Pawnee Montane Skipper  
(Hesperia leonardus montana)**

Federal status: Threatened  
Colorado status: Not listed

The upper side of this butterfly's wing is brownish red with distinct yellowish spots near the outer margins. The underside is brown with cream spots. The wing-span is from 1.5 to 2 inches.

Habitat: This species is found in dry, open, Ponderosa pine woodlands at elevations from 6,000 to 7,500 feet. The slopes are moderately steep with soils derived from Pikes Peak granite. The understory is limited in the pine woodlands. Blue Grama grass is the larval food plant, and the prairie gayfeather is the primary nectar plant. The species is found in Douglas, Jefferson, Park and Teller counties. The total known habitat within the range is estimated to be 37.9 square miles.

